



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

ently does not belong to that genus at all, since it is without the very striking internal transfusion sheath of the lower cretaceous genus, has a double instead of a single foliar bundle (a point of great importance, as will be recognized by those versed in the anatomy of *Pinus* in its living species), and no centripetal xylem. The authors make light of the absence of the last feature, but in this they are apparently not well advised.—E. C. JEFFREY.

**Conduction of stimulus.**—ROTHERT has shown that the conduction of stimulus of unilateral illumination from the tip of the *Avena* seedling to the darkened basal portion occurs when the vascular strands are cut; and that a horizontal incision on one side, whether toward, away, or on the flank in reference to the one-sided illumination, still permits conduction. FITTING showed that when the incision was away from the light and a mica plate inserted, no conduction occurred. The insertion of a slice of rattan in the same position did not prevent conduction; the latter of course allows the continuity of water and solution. When the mica plate was inserted in an incision on the lighted side, conduction was not hindered. JENSEN<sup>14</sup> finds that with the incision away from the light, no conduction occurs in dry air or in water. He assumes that under favorable conditions the stimulus can be conducted across the wound while under unfavorable conditions it cannot. In saturated air the stimulus was also conducted from the tip to the darkened base, even after the tip (1 cm. long) had been entirely cut off and set back and fastened by gelatin and cocoa butter.—WILLIAM CROCKER.

**Embryo sac of Pandanus.**—In 1909 CAMPBELL published an account of the embryo sac of *Pandanus*, which was reviewed in this journal (47:485, 1909). In this Javanese material the fertilization stage was not secured, so that it was not certain that the interesting situation described is the one at fertilization. Now there has come to hand additional material (*P. coronatus*) which has supplied the missing stage.<sup>15</sup> An ordinary egg apparatus is organized, but there occurs "a large discoidal mass of cells" at the antipodal end of the sac, and fusions of "polar" nuclei (up to six) were observed. The number of cells in the sac at the fertilization stage would thus seem to be greater than that recorded for any other angiosperm. The amount of antipodal tissue suggests the situation in *Sparganium*, the difference being that in the latter genus this tissue develops after fertilization.—J. M. C.

<sup>14</sup> JENSEN, P. BOYSEN, Ueber die Leitung des phototropischen Reizes in Avenakumpflanzen. Ber. Bot. Gesell. 28:118-120. 1910.

<sup>15</sup> CAMPBELL, DOUGLAS H., The embryo sac of *Pandanus coronatus*. Bull. Torr. Bot. Club 24:293-295. figs. 6. 1910.